

One Earth

Review

Ocean conservation boosts climate change mitigation and adaptation

Juliette Jacquemont,^{1,2,3} Robert Blasiak,^{4,5} Chloé Le Cam,² Maël Le Gouellec,² and Joachim Claudet^{1,*}

¹National Center for Scientific Research, PSL Université Paris, CRIOBE, CNRS-EPHE-UPVD, Maison de l'Océan, 195 rue Saint-Jacques, 75005 Paris, France

²Ocean & Climate Platform, 195 rue Saint-Jacques, 75005 Paris, France

³School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195, USA

⁴Stockholm Resilience Centre, Stockholm University, 106 91 Stockholm, Sweden

⁵Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

*Correspondence: joachim.claudet@cnrs.fr

https://doi.org/10.1016/j.oneear.2022.09.002

SUMMARY

Marine protected areas (MPAs) are increasingly being promoted as an ocean-based climate solution. However, such claims remain controversial because of the diffuse and poorly synthesized literature on climate benefits of MPAs. To address this knowledge gap, we conducted a systematic literature review of 22,403 publications spanning 241 MPAs and analyzed these across 16 ecological and social pathways through which MPAs could contribute to climate change mitigation and adaptation. Our meta-analysis demonstrates that marine conservation can significantly enhance carbon sequestration, coastal protection, biodiversity, and the reproductive capacity of marine organisms as well as fishers' catch and income. Most of these benefits are only achieved in fully or highly protected areas and increase with MPA age. Although MPAs alone cannot offset all climate change impacts, they are a useful tool for climate change mitigation and adaptation of social-ecological systems.

INTRODUCTION

Climate change has started to undermine human well-being and planetary health, generating a sense of urgency to identify effective mitigation and adaptation strategies.¹ Recent years have seen a growing focus on the ocean's central role in climate² as well as an increase in associated advocacy efforts. This is reflected, among other things, in the increasing inclusion of ocean issues in nationally determined contributions for climate mitigation and adaptation.³

Considerable consensus exists on the cascading impacts of climate change on marine ecosystems and coastal communities,^{2,4-6} but no such consensus exists on ocean-based climate solutions.⁷ Most notably, although the potential of ocean conservation tools such as marine protected areas (MPAs) is widely recognized to deliver multiple positive ecological and social outcomes,^{8,9} their ability to contribute to the resilience of marine social-ecological systems to climate change or to contribute to carbon sequestration remains highly controversial.^{10–13} Although dogmatism drives part of these disagreements,¹⁴ the underlying scientific challenge lies in the diverse and diffuse body of research that allows us to formulate a range of hypotheses on whether marine conservation can serve as a climate solution. Some argue that the increases in abundance, biomass, and biodiversity of fished populations occurring in MPAs^{15–17} promote other ecological outcomes, such as reproductive output, genetic diversity, or ecosystem stability,¹⁰ ultimately contributing to the adaptive potential of marine ecosystems to climate change.¹⁸⁻²

However, few studies have directly tested the effects of MPAs on climate change adaptative potential, and the existing literature shows contrasting results.^{21,22} A great amount of literature has shown that MPAs failed to protect coral reefs from bleaching during heat waves.²³ In some instances, coral loss has even been shown to be greater in MPAs than in unprotected areas. This phenomenon, referred to as the "protection paradox,"¹² has fueled much of the opposition in advocating MPAs as a tool for climate adaptation.

The effects of marine conservation on social adaptive potential are another contentious topic because millions of livelihoods directly depend on fisheries for income and food security. Although MPAs have been shown to provide benefits to fisheries because of spillover of fish from protected to fished areas,²⁴ controversy remains regarding the overall costs and benefits to fishing communities.²⁵⁻²⁷ Because coastal populations are among the most vulnerable to climate change,^{1,2} it is paramount to identify solutions that will not further negatively impact their adaptive capacity.^{28,29} However, no synthesis of the literature examining how MPAs impact social adaptive potential exists, casting doubt on whether marine conservation can be used as a holistic tool that benefits marine ecosystems and coastal communities.³⁰

The debate has also centered on the ability of MPAs to provide climate mitigation benefits, notably through increased carbon sequestration from rebuilt fish populations or from undisturbed sediments from trawling bans.^{13,31} Clarity regarding how ocean conservation contributes to ecological adaptation, social adaptation, and climate change mitigation is urgently needed to



ensure that effective ocean-based strategies are adopted in climate change policies.

To address these knowledge gaps, we identified the potential pathways through which MPAs could contribute to climate change mitigation and adaptation (hereafter referred to as "climate pathways"). A layered typology was used to classify these climate pathways: first along mitigation and adaptation dimensions and then distinguishing among ecological and social dimensions of adaptation. We then carried out a systematic literature review and summarized the results of all empirical studies documenting MPA outcomes on these climate pathways using vote counting and a meta-analytical approach. Vote counting (i.e., calculating the fraction of studies reporting positive, negative, or neutral outcomes) allows us to synthesize results from both qualitative and quantitative studies and to overcome some publication biases associated with meta-analysis. For these reasons, it is commonly used in social science,^{25,32} as well as in ecology.^{33,34} When sufficient quantitative data were available to perform meta-analysis, we also quantified the direction, magnitude, and uncertainty of MPA outcomes on climate pathways. Because previous analyses have shown that high levels of protection are required to achieve ecological¹⁵ and social³⁵ outcomes, we also investigated whether this was a necessary condition for MPAs to produce climate mitigation and adaptation benefits.

Our literature review found that marine conservation enhances most ecological and social climate pathways. Meta-analyses showed that MPAs significantly increase carbon sequestration, coastal protection, biodiversity, and the reproductive capacity of marine organisms as well as fishers' catch and income. However, these benefits were only achieved under full or high levels of protection. Our study provides evidence that MPAs constitute an effective solution for climate change mitigation and adaptation of the intertwined components of social-ecological systems.

MPA MITIGATION AND ADAPTATION PATHWAYS

Sixteen pathways through which MPAs could contribute to climate change mitigation and adaptation were identified by drawing on reviews of social and ecological outcomes of MPAs^{10,12,20,25,28} (Table S1). Two climate pathways contributed to climate mitigation (carbon sequestration and local acidity buffering) and 14 to climate adaptation (Figure 1). Social adaptation pathways were derived from the five pillars of social adaptive capacity²⁸: assets, flexibility, agency, learning, and social organization, to which was added food security.³⁰ Ecological pathways were derived from climate adaptation pathways described previously:10,12,20 connectivity, phenotypic plasticity, genetic diversity, biodiversity, stability, reproductive output, body condition, and coastal protection. Up to two indicators were selected to measure each pathway based on the most common indicators used in the reviewed studies. Additional indicators were also investigated when they allowed us to quantify aspects of the studied pathways not yet captured by the first two indicators (Table S2; Figure S4). The definition of each pathway and the units used to measure each indicator are detailed in Table S2. For two pathways (carbon sequestration and coastal protection), we also included all studies comparing exploited and preserved marine areas even when not officially labeled MPAs. This was done

because the literature documenting the effect of preservation initiatives, comparable with MPAs, is abundant, whereas almost no study directly documenting MPAs is available.

The systematic literature review on MPA effects on climate pathways generated a total of 22,403 publications, of which 378 were included in vote counting, providing insights from 241 different MPAs. Publications were unevenly distributed among continents (Figures 1B–1D), with most ecological adaptation pathways studied in Europe and most mitigation and social adaptation pathways studied in Asia (Figure S2). We found empirical evidence documenting the effects of MPAs on all climate pathways except acidity buffering, connectivity, and phenotypic plasticity (Figure 1). Eight climate pathways had sufficient quantitative data (n > 3) to perform a meta-analysis.

MARINE CONSERVATION CONTRIBUTES TO CARBON SEQUESTRATION

We investigated the effects of marine conservation on the carbon (C) sequestration capacity of six marine C sinks: three blue C ecosystems (mangrove, tidal marshes, and seagrass), which have already been recognized by the Intergovernmental Panel on Climate Change (IPCC) for C accounting schemes,³⁶ as well as sediments, macroalgae, and fish.^{31,37} C sequestration was defined as organic C stored for over 100 years.³⁸ Mean effect sizes (log response ratios [InRR], 95% confidence interval) indicated significant increases in C sequestration in preserved or restored seagrass ($InRR = 0.76 \pm 0.34$) and mangrove (InRR = 0.75 ± 0.14) ecosystems in comparison with similar areas undergoing human pressure (e.g., thinning, anchoring, conversion to plantations). Similarly, sediments in untrawled seabed sequestered significantly more C than areas exposed to trawling (InRR = 0.13 ± 0.10). Conservation had no effect on C sequestered by tidal marshes (Figure 2D), mostly because conversion of unprotected marsh into agricultural land increased C sequestered in plant biomass and in the soil.³⁹ Partial (e.g., thinning, anchoring) or full (e.g., clear cutting, excavating) degradation of mangroves and seagrass resulted in similar decreases of sequestered C, indicating that even low levels of human impact result in important C emissions (Figure S6). No studies documented the effect of protection on the quantity of sequestered C originating from fish or macroalgae biomass. However, many studies have shown that MPAs significantly increase fish biomass (lnRR = 1.10 ± 0.58),¹⁶ which can serve as a proxy for C sequestration because a portion of that biomass undergoes exportation and is subsequently sequestered in the deep sea.⁴⁰ This is supported by several studies that calculated large impacts of fisheries on C sequestration from fish biomass removal.^{31,41} However, this remains an indirect measurement of C sequestration, and more research is needed to quantify the exportation rates of organic C from fish biomass toward sediments.⁴² Although macroalgae are increasingly being advocated as significant actors in marine C sequestration,^{37,43} major knowledge gaps remain regarding how marine conservation affects their living biomass and, thus, their contribution to C sequestration.

Although the effect of MPAs on local acidity buffering was not documented by any study, our meta-analysis revealed that seagrass increased mean local pH and that mangroves and macroalgae decreased it (Figure 3). Benefits to the adaptive potential of





Figure 1. Pathways through which MPAs can contribute to climate change mitigation or adaptation, with quantitative and spatial assessment of available information

(A) List of the climate pathways identified and their associated indicators. Blue, green, and orange color themes highlight the classification of pathways among mitigation, ecological adaptation, and social adaptation categories, respectively. Social and ecological adaptation pathways are shown linked to the components of adaptation (i.e., resistance, recovery, and adaptive potential) to which they contribute. The number of articles found through our search queries and the number of articles that met the screening criteria for vote counting and meta-analysis are indicated for each pathway. The width of bars is proportional to the number of studies selected at each step. CPUE stands for catch per unit effort.

(B–D) Location of the study sites documenting pathways of climate mitigation (B), ecological adaptation (C), and social adaptation (D). The size of dots is proportional to the number of studies at a given location for a given pathway.

marine organisms could arise from exposure to greater pH fluctuations that occur in vegetated habitats, which has been shown to increase tolerance to acidification.⁴⁴ More research is required to support this hypothesis and assess whether conservation of vegetated habitats could benefit pH buffering or acidity tolerance.

CONTRIBUTION OF MPAs TO ECOLOGICAL ADAPTATION

We found that MPAs contribute to ecological adaptation by increasing biodiversity, reproductive output, and coastal protection compared with unprotected sites (Figure 2C). Species richness was higher in MPAs (lnRR = 0.20 ± 0.10), in agreement with previous regional¹⁷ and global⁴⁵ meta-analyses, but the Shannon index was unchanged between protected and unprotected sites (lnRR = 0.06 ± 0.06). Increased species richness could play a central role in climate adaptation because more species increase the odds that ecosystem functions are maintained even after a stressor eliminates certain species,⁴⁶ a concept referred to as the insurance hypothesis.⁴⁷ Effects of MPAs on

the reproductive output of marine organisms (measured as increased larval densities and egg production) had the greatest magnitude of all studied pathways (lnRR = 1.21 ± 0.96). This likely results from larger, older, and more abundant individuals in MPAs, leading to increased production of offspring.⁴⁸ In contrast, recruitment rates showed no significant increase, which could be explained by the higher predation rates on recruits experienced in MPAs.⁴⁹ Reproductive capacity is an important attribute of adaptation because it is linked to faster rates of population recovery⁵⁰ and increased larval dispersal distance, which, in turn, can promote populations' connectivity and ability to colonize new habitats.⁴⁸

Preservation of mangroves and tidal marshes enhanced coastal protection through soil accretion rates of an additional 1.38 ± 0.88 cm year⁻¹ compared with degraded habitats. Coral reefs and seagrass can also contribute to coastal protection through wave attenuation (Figure 3), but no studies have assessed whether wave attenuation is enhanced when habitats are in an MPA. However, wave attenuation is linked to a habitat's structural complexity and vegetation density,^{51,52} which suggests that habitats protected from physical disturbance would

One Earth

Review







Figure 2. Effects of marine protected areas (MPAs) on climate change mitigation and adaptation pathways

(A–C) Climate pathways (A) and direction of reported MPA effects from studies included in the vote-counting analysis (B) and magnitude of outcomes from studies included in the meta-analysis (C). In (B), the x axis indicates the cumulative number of studies reporting positive outcomes (right side of the bar plot, green) and ambiguous, negative, or neutral outcomes (left side of the bar plot). In (C), the x axis indicates the log-transformed ratio of indicators between MPAs and controls. In the case of coastal protection, the effect size was calculated as the Euclidian difference between MPAs and controls, hence the separate scale provided. Values are presented as mean values ± 95% confidence interval. Black dots indicate effect sizes that do not overlap zero and white dots those that overlap zero. Sample sizes (i.e., number of studies) of vote counting and meta-analysis results are indicated by n values. *In the case of C sequestration from fish biomass, the effect size was calculated from three previous meta-analyses, representing data from many more individual studies.

(D–F) Effect of habitat type (D), protection level (E), and presence of a fully protected area (F) on the capacity of MPAs to provide climate benefits. Colors, sample size, and x axis have the same meaning as in (C). †Effects of full protection on C sequestration (fish biomass only) are reported from Lester et al.¹⁶ ‡Effects of high and low protection on C sequestration (fish biomass only) are reported from Lester et al.¹⁵

perform better than degraded habitats. We found that MPAs increased ecosystem stability, but quantitative evidence remains too low to draw general conclusions (lnRR = 0.10 [-0.03, 0.63], n = 5). MPAs had no effect on genetic diversity (measured as allelic richness, expected heterozygosity, and observed heterozygosity; Figures 2C and S7). This is consistent with the one previous regional meta-analysis⁵³ that also found no effect of MPAs on genetic diversity.

Body condition could only be assessed through vote counting, and studies mostly reported a neutral effect of MPAs (Figure 2B). Connectivity and phenotypic plasticity were not addressed by any studies. Greater larval and egg densities in MPAs do indicate an increased potential for dispersal, but whether this potential effectively translates into increased connectivity with adjacent areas remains to be evidenced, using indicators such as fixation index.⁵⁴ The ability of MPAs to enhance connectivity constitutes



a major knowledge gap because connectivity is one of the most important pathways for adaptation to climate change by allowing recolonization of disturbed habitats, gene exchange, and climatic migration.⁵⁵

CONTRIBUTION OF MPAs TO SOCIAL ADAPTATION

CelPress

Among social adaptation pathways, food security and assets displayed sufficient quantitative data to perform a meta-analysis. MPAs significantly increased food security ($InRR = 0.43 \pm 0.21$; Figure 2C), measured using catch per unit effort (CPUE). Increases in CPUE arise from replenishment of fish populations in MPAs that can subsequently spill over into fishing grounds, benefitting local fisheries.²⁴ Increases in CPUE occurred when comparing fishing in and outside of MPAs, in the case of partially protected areas, and when comparing fishing near and far away from MPAs (Figure S8). In both cases, we only observed increases in CPUE when the MPA included or was a fully protected area; i.e., a protected area where no fishing is allowed (Figure 2F). Our results oppose those of a regional meta-analysis⁵⁶ that found a negative impact of MPAs on the CPUE of southern European fisheries. We suggest that our results, built on a higher number of studies and a wider geographical scale, overcome the specificities of this previous meta-analysis and better reflect the effects of MPAs on fisheries. This result is backed up by the observed increase in fishers' income in the presence of a fully protected area ($InRR = 0.35 \pm 0.29$). Increased fishers' income demonstrates that MPAs can potentially offset costs generated by displacement of fishing grounds. Increases in CPUE and income support the long-running controversy that MPAs can enhance the livelihood of fishers, which, in turn, con-

Figure 3. Contribution of marine habitats to climate pathways

Values in barplots indicate the capacity of marine habitats to contribute to C sequestration (mg C km⁻²), coastal protection (accretion in mm year⁻¹ and wave attenuation in % wave decrease), and acidity buffering (Δ pH unit). The height of the barplots represents the capacity of habitats to contribute to a given indicator relative to the maximum value reached by a habitat. For pH buffering, negative values indicate that the habitat decreases the average local pH. The spatial extent of habitats is indicated under each habitat's name and is represented by a half-sphere area that is log proportional to its global spatial area.

tributes to building social adaptive capacity to climate change through increased food security and assets.

MPAs had a majority of positive outcomes across all social adaptive pathways (i.e., 8 indicators out of 11) with the exception of social organization (Figures 2B, S3, and S4). In addition to CPUE and income benefits, the most positive social adaptive outcomes of MPAs were increases in environmental awareness (95% of cases), participation (57% of cases), and alternative livelihood (48% of cases). These findings

are consistent with previous systematic reviews.^{25,32} The three indicators for which we found negative outcomes of MPAs were user rights, conflict, and costs (used to measure agency, social organization, and assets pathways, respectively). In 65% of cases, user rights were negatively affected by access and extraction restrictions imposed by MPA regulation, reducing local people's agency.^{27,57} Increases in conflicts (59% of cases) arose from changes in spatial use of marine areas, potentially leading to tensions between different activity sectors; e.g., tourism operators and fishers.⁵⁸ However, MPAs increased social cohesion (56% of cases), likely resulting from a shared conservation vision and sense of purpose between stakeholders.⁵⁹ Such a discrepancy of MPA outcomes on conflict and cohesion suggests that impacts of MPAs on social organization are strongly context dependent.

CONDITIONS UNDER WHICH MPAs DELIVER CLIMATE BENEFITS

We found sufficient data to investigate the effect of the level of protection, age, and size of MPAs on four climate pathways (C sequestration, biodiversity, CPUE, and income).

Across all four pathways, only full and high levels of protection resulted in mitigation or adaptation benefits (Figure 2E). In contrast, lower levels of protection generated no benefits. Increases in species richness and in fishers' income only occurred for fully protected areas, where no fishing is allowed. Increases in CPUE and C sequestration were also achieved by highly protected areas, where low-impact fishing is allowed. However, this was only the case when a fully protected area was also present in the MPA considered (i.e., a multi-zone MPA with one zone being a fully protected area).



Figure 4. Disentangling confounding factors of the protection paradox

Effect sizes were calculated as natural log ratios of MPA and controls for resistance, recovery, and expected ecological adaptation of marine organisms. Values represent mean ± 95% CI. Black effect sizes indicate CIs that do not overlap zero, and white effect sizes indicate CIs that do overlap zero. Colored dots represent individual effect sizes calculated from each publication that were included in the overall effect size represented. Colors of dots represent the taxa of organisms studied in each publication. Donut diagrams on the right represent the distribution of individual effect sizes among taxa. Contributing pathways indicate the pathways from which individual effect sizes were aggregated to obtain the effect size of expected ecological adaptation.

All four climate pathways were positively correlated with MPA age (Figure S9; Table S5). This is explained by the time required for exploited fish stocks to rebuild and subsequently benefit adjacent fisheries. Similarly, C sequestration was positively correlated with the number of years since restoration of mangroves and seagrass. CPUE was positively correlated with MPA size (Figure S10), which can result from a greater proportion of mobile fish having their home ranges included in the MPA.⁶⁰ Level of protection, age, and size have already been established as important drivers and enabling conditions for MPAs to deliver positive ecological outcomes.^{15,61} Our results show that this is also the case for climate mitigation and adaptation benefits.

In addition to size, age, and protection level, the magnitude of climate benefits achieved through protection depends on the marine habitat considered. The characteristics of each marine habitat influence its ability to contribute to climate pathways, and its spatial extent determines its maximum potential contribution to climate change adaptation and mitigation (Table S6). This is particularly true for the C sequestration, coastal protection, and acidity buffering pathways, which derive from ecosystem services delivered by specific habitats. Habitats with high C sequestration potential but low spatial extent (e.g., mangroves and seagrass; Figure 3) translate into a high mitigation potential per surface area but low mitigation potential at the global scale. Conversely, fish and sediments have lower mitigation potential per surface area but can play an important role in global climate mitigation because of their vast extent. More research on sediments and new or updated IPCC guidelines for including sediments and other marine C pools into national greenhouse gas accounting would help make MPAs more actionable in climate change mitigation and including them into nationally determined contributions (NDCs).

FROM ADAPTATION PATHWAYS TO RESILIENCE

To test whether the enhanced ecological pathways found in this study effectively translate into increased climate adaptation, we performed two complementary meta-analyses quantifying the effect of MPAs on the resistance and recovery of marine organisms to climatic stressors. Our systematic literature review identified 19 papers that focused exclusively on warm-water coral species. We found that MPAs had a negative effect on the resistance of corals to climatic stressors (InRR = -0.38 ± 0.36) and no effect on their recovery from climatic stressors (InRR = $0.04 \pm$ 0.23) (Figure 4), which is consistent with previous findings²³ and referred to as the "protection paradox."¹² In contrast, we found a significantly positive effect of MPAs on expected ecological adaptation (InRR = 0.31 (0.03-0.87)), calculated from all data points included in resilience-related pathways (i.e., stability, biodiversity, genetic diversity, and reproduction). The gap between the expected adaptation benefits that should emerge from enhanced ecological pathways and the observed absence of improvement in resistance and recovery of marine organisms could result from (1) the fact that the adaptation pathways we measured are inappropriate or too indirect to inform on adaptation capacity or (2) differences in the studied system between recovery/resistance studies and adaptation pathways studies. The latter hypothesis likely plays an important role in this gap because all studies on recovery and resistance focused on corals, whereas most studies documenting adaptation pathways focused on fished species. Not only is the literature on conservation outcomes on resilience biased toward coral species, but it has also been shown to be further biased toward specific oceanic regions.⁶² The ability of MPAs to enhance ecological adaptation results mostly from alleviation of fishing pressure, which represents a minor threat to corals in comparison with temperature rise and acidification.⁶³ In contrast, fishing pressure is currently the main threat to fished species,⁶⁴ which is why fully protected areas were observed to enhance several adaptation pathways. As a result, the "protection paradox" observed for corals cannot be generalized to other taxa, and dedicated studies should be performed to test how MPAs affect the climatic resilience of fished species. Although evidence remains scarce, theoretical^{65,66} and empirical^{22,67} research suggests positive effects of MPAs on the resilience of fish populations. More research is required to test whether the enhanced adaptation pathways reported in this study effectively translate into enhanced resilience of fished species.



CLIMATE-SMART MPA DESIGNS

A growing body of literature has focused on the design characteristics of "climate-smart" MPA features, aimed at maintaining current conservation benefits under future climatic conditions. Recommendations have included MPA networks that account for future patterns of connectivity under changing oceanic currents, spatially dynamic MPAs that follow species migrations, or protection of climatic refuges such as upwelling sites.68,69 Although these elements are certainly worth considering when designing an MPA, a recurring theme in our results is that MPAs must be highly or fully protected to provide a broad range of benefits, including positive climate change mitigation and adaptation outcomes. The clear signal that level of protection should take precedence in MPA design is in stark contrast to the current paradigm of ocean protection globally, with the proportion of fully and highly protected MPAs worldwide plummeting as countries have rushed to meet conservation commitments.70-72 Ensuring high protection levels in existing MPAs should be the priority to secure climate benefits to coastal social-ecological systems.

MPA GOVERNANCE

The diverse effects of MPAs on social organization and agency of coastal communities underscore that governance is a fundamental factor in determining whether MPAs will enhance or impede social adaptation. The outcomes of several social adaptation pathways, such as agency and learning, result from how local stakeholders are included throughout MPA establishment and management.^{27,73} Involvement of stakeholders through co-management allows the identification of peoples' needs and minimization of negative social impacts, e.g., through the development of alternative livelihood options or conflict-resolution tools.⁷⁴ Positive perceptions of governance by local stakeholders has also been shown to increase support and compliance with MPA regulations,⁷⁵ which, in turn, allows ecological benefits to accrue.⁷⁶ This results in a positive feedback loop whereby perceived ecological benefits act as another incentive for communities to support MPAs⁷⁷ and allow food security and asset benefits to accrue. To promote social adaptive capacity, there is benefit for MPA managers to thoroughly consult and inform local communities about MPAs' goals and rules;⁷⁸ involve local communities in MPA management and decision-making;⁷⁹ identify potential negative social outcomes of MPAs and find solutions to avoid, reduce, or compensate for them;⁸⁰ and monitor MPA social outcomes and adjust MPA governance accordingly.73

WHAT ROLE FOR OCEAN CONSERVATION IN CLIMATE POLICY?

Understanding the benefits and potential of effective ocean conservation has taken on particular urgency in international policy. Appealing narratives of ocean conservation have found resonance in civil society as well as governments, resulting in calls to protect 30% of the ocean by 2030 and expectations of an imminent 30% target from the Convention on Biological Diversity. Simultaneously, "blue carbon" projects have become an attractive pros-

One Earth Review

pect for the private sector, with a 50-fold growth in the blue carbon offset market projected by 2030.⁸¹ When carefully navigated, such targets and trajectories represent a window of opportunity to underscore the benefits of ocean conservation for nature, the climate, and human well-being. Likewise, an incautious or non-inclusive approach could lead to large-scale dispossession of local communities from the landscapes and seascapes that sustain them,⁸² and poorly implemented protected areas could fail to deliver the expected benefits, undermining perceptions of MPAs as an ocean-based climate solution.

Our synthesis provides empirical evidence that MPAs contribute to climate change mitigation and adaptation, particularly when fully protected, making them a key tool for achieving the goals of the United Nations Sustainable Development Agenda and the Paris Climate Agreement. Multiple opportunities now exist to leverage this scientific basis to guide public policy and private sector initiatives. First, there is considerable scope to expand the consideration and recognition of MPAs in national climate strategies, including NDCs and adaptation communications.⁸³ Such efforts would be aided by further allocation of resources to accelerate the expansion of IPCC-recognized blue carbon strategies to include marine sediments, enabling reliable accounting for mitigation benefits of MPAs with trawling restriction.³⁶ Second, because 64% of the ocean falls outside of national jurisdictions, it is also fundamental that ongoing negotiations on a treaty for the high seas are successfully concluded and enable designation of area-based management tools such as MPAs. Finally, the sense of urgency to act and scale up innovative solutions should be welcomed, but not at the expense of recognized best practices regarding inclusive MPA governance and high to full levels of protection.

EXPERIMENTAL PROCEDURES

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Joachim Claudet (joachim.claudet@cnrs.fr).

Materials availability

This study did not generate new unique materials.

Data and code availability

All original code and datasets have been deposited at Zenodo under https:// doi.org/10.5281/zenodo.7108799 and are publicly available as of the date of publication.

Identification of pathways

Pathways through which MPAs could contribute to climate change mitigation or adaptation were identified from the literature. Review articles on the ecological and social adaptive capacity of marine systems were screened, and the components of marine climate change mitigation and adaptation listed therein were attributed to climate pathways. Five reviews were retained, ^{1012,20,25,28} from which a total of 16 pathways were identified, as detailed in Table S1. A description of each pathway identified (definition, indicator, unit) is detailed in Table S2. Definitions of social adaptation pathways were adapted from Cinner et al.²⁸

Systematic literature review search strategy

To investigate the 16 pathways identified in the section "identification of pathways," nine independent systematic searches of published peer-reviewed literature were performed on Web of Science. Social adaptive pathways were reviewed through a unique systematic search because most social studies tackled several social pathways, making it more effective to screen articles for all social pathways at once. Following the same logic, the pathways "body condition," "phenotypic plasticity," "resistance," and "recovery" were reviewed jointly, as well as "connectivity" and "reproductive potential." All other pathways were reviewed individually.



To make sure no important article was missed, the database obtained through our systematic literature searches was complemented with studies included in previous systematic reviews relevant to one of our pathways. The following systematic reviews were used to complete our database:

- For social pathways²⁵
- For C sequestration^{13,84,85}
- For resistance and recovery²³

Additional articles identified from expert knowledge or from citations in reviewed publications were also added to our database.

All systematic reviews were performed using Web of Science with no time limitation and using the "topic" (titles, abstracts, and keywords) search section. The dates when the searches were performed are detailed in Data S1 (accessible through the Zenodo link provided in the "data and code availability" section). All search strings were composed of two parts: a part describing the system of interest and a part describing the climate pathway. The full search string used for each systematic review can be found in Data S1. Search strings were validated by making sure a set of five pre-selected studies that matched our search criteria were found.

For adaptation pathways, the system of interest was limited to MPAs, and the search string included all different designations used to refer to MPAs in the literature: (("locally managed marine area*") OR ("locally-managed marine area*") OR ("community-based marine area*") OR ("marine communitybased area*") OR ("fisher* closure*") OR ("marine closure*") OR ("marine restricted area*") OR ("fisher* closure*") OR ("marine closure*") OR ("marine restricted area*") OR ("restricted marine area*") OR ("taboos") OR ("tabus") OR ("rahui") OR ("marine protected area*") OR ("marine reserve*") OR ("marine reservation area*") OR ("marine sanctuar*") OR ("marine conserved area*") OR ("marine conserved territor*") OR ("ocean conservation area*") OR ("marine community-based conservation area*")).

For mitigation pathways and coastal protection, the system of interest was enlarged to preserved habitats because mangroves and marshes benefit from a variety of conservation measures that act similarly as MPAs by regulating natural resource extraction. To allow a maximum of results to be found, the search string broadly included the names of the habitats of interest (Data S1; see "data and code availability" section), and the articles were selected when they compared preserved habitats with degraded controls (see "screening criteria" section).

Screening criteria for the vote-counting approach

Screening of articles was performed using a three-stage process: screening of titles, screening of abstracts, and screening of full texts. Some criteria were common to all systematic reviews, and others were specific to pathways. Common screening criteria were as follows:

- The article provided primary data collected on the field or in a lab (models, reviews, and syntheses were excluded, as well as studies using datasets already presented in previous studies).
- The system studied was a marine or coastal habitat (freshwater systems were excluded).
- The study compared an MPA site with a non-MPA site (or a preserved/ restored habitat with a degraded habitat in the case of mitigation pathways and coastal protection), using a before-after or control-impact design (before-after-control-impact designs were extracted when available).
- Studies with pseudo-replication were excluded.

In the case of studies comparing MPAs with control sites (all studies documenting adaptation pathways except coastal protection), screening criteria were as follows:

- The control site must be outside of the MPA (comparisons between notake zones and buffer zones or partially protected areas were excluded).
- The MPA must be a permanent closure (seasonal fishery closures were excluded).

Screening criteria specific to each pathway are detailed in Table S2.

Title and abstract screening were conservative to avoid excluding any relevant studies and were used to phase out studies that were clearly irrelevant to our search. Studies that passed our title screening phase were exported from Web of Sceince in an Excel file with all their relative information (including abstracts). Abstracts of the selected articles were then screened using the abstract screener interface provided by the {metagear} package in R Core Team (2021).⁸⁶ This package allows us to screen abstracts without showing authors or the journal's title, limiting sources of bias in the selection process. Articles selected after abstract screening were downloaded, and their text was fully read to assess whether they were relevant to our search criteria.

Selection of indicators for each pathway

A list of indicators used to describe pathways was identified from the studies that met all screening criteria. When different indicators informed on a same aspect of a pathway, they were grouped together (e.g., larval density and egg density were grouped under reproductive output). When different indicators informed on different aspects of a pathway, they were kept distinct (e.g., reproductive output and recruitment for the "reproduction" pathway). The list of indicators selected for each pathway is detailed in Table S2. For clarity of figures and content, a maximum of two indicators per pathways were presented in the main article. Indicators presented in the main article were chosen to represent contrasting MPA outcomes (e.g., a positive and a negative outcome within the same pathway).

Number of values extracted per study

In the following cases, several values were extracted from a same article:

- When different MPAs were studied.
- When different ecological groups (fish, fished invertebrates, corals, algae, seagrass ...) were studied.
- When different stakeholder groups (fishery, tourism, recreational, and coastal community) were studied.
- When MPA zones with different levels of protection were studied.

In all other cases, values were averaged between categories (e.g., different species of the same taxon, different religions within a community ...) to avoid over-representing studies. When time series of values were presented for a given MPA, only the most recent data point was kept as the MPA "after" value, and the data point closest to before MPA establishment was kept as the MPA "before" value.

Qualitative data extracted from screened articles

For each value extracted, the following information was also extracted:

- Design of the study: before-after (BA), control-impact (CI) or BACI. For social studies, the category "perception" when MPA outcomes were based on interviews and stakeholders' perceptions.
- MPA name, MPA size (km²), and MPA age (year) at the time of data collection. MPA age was counted from the moment when the MPA was enforced rather than announced to account for years during which marine ecosystems were effectively protected. When MPAs underwent several stages of extension, we counted MPA age from when the latest extension was carried out. When MPA features were not reported in the publication, we used information from the MPA atlas (https://mpatlas.org/).
- The level of protection of the zone studied by the MPA and, when present, the size of the fully protected area in the MPA studied (km²). Levels of protection were determined according to the regulation-based classification system for MPAs.⁸⁷ In addition to information provided in the study, complementary searches were performed using the MPA atlas and the official management declarations of MPAs to determine the level of protection.
- Continent, oceanic region, climate of the study location.
- Ecosystem, taxon, and species studied.
- Type of governance and of stakeholder studied for social pathways.
- Pathway, indicator, unit, and type of error provided in the study.

One spatial point was mapped for each data point used in vote counting and meta-analysis. Spatial points were located at the centroid of the protected area, as found in the MPA atlas (https://mpatlas.org/). Mapping was carried out using the {sf} package to manipulate vectors, {maps} to create the back-ground layer, and {tidyr} to manipulate datasets. The projection used was Mollweide.

Direction of MPA outcomes reported in vote counting

For each article meeting our search criteria, the direction reported by the article (i.e., how marine protection influenced the outcomes of the pathway) was extracted. The direction of outcomes was positive, negative, neutral, or ambiguous. "Ambiguous" was used when contradicting trends were reported by the article and when the article could not conclude how the MPA affected the pathway. Increases in positive indicators (e.g., participation, alternative livelihoods, and catch) were coded as positive (and negative for decreases), and increases in negative indicators (conflict, costs, mortality, coefficient of variation) were coded as negative for decreases).

When results were quantitative and associated with an error, the direction "neutral" was attributed when the 95% confidence interval overlapped zero. When results were qualitative or when no error was associated with the results,



the direction of the article was determined using the authors' interpretation of the results in the article's text. In social studies, the directions of outcomes were often given as percentages, in which case the prevalent category was chosen for the global direction of the study (e.g., 45% positive, 25% neutral, and 35% negative were reported as positive). When two outcomes' proportions were closer than 5% to each other (e.g., 48% negative and 52% positive), "ambiguous" was attributed.

Additional screening criteria for the meta-analyses

Among studies that met our search criteria for vote counting, we identified those that provided standard statistical data required to perform a meta-analysis; i.e., mean, error, and sample size. Studies providing median values instead of means were excluded. We subsequently identified the most common unit to measure a given indicator in the selected studies and excluded all studies that used units that were not comparable. Several indicators were kept for the same pathway when they informed on different aspect of that pathway (e.g., species richness and Shannon index for biodiversity). The list of indicators kept for each pathway is detailed in Table S2. A meta-analysis was only performed when more than three articles met our criteria for a given indicator.

Quantitative data extraction and preparation

Quantitative results were extracted from the main text, tables, and figures of articles and from supplemental material. Mean values and errors were extracted from figures using the online application WebPlotDigitizer v.4.0 (Rohatgi 2018). Data extraction of the 16 pathways was carried out by three of the co-authors of this study. Data extraction of five studies was first carried out by all co-authors to make sure that the methodology was acquired by all. All data extracted by a co-author were double-checked by another coauthor to reduce extraction or calculation mistakes.

All types of errors (standard errors, variance, confidence intervals) were converted into standard deviation (SD). Mean values and variance were calculated when results were presented across categories irrelevant to our study (see "number of values extracted per study" section), and sample size was added. The same was done when a publication presented multiple control sites for a unique MPA site. The variance of mean values s²_{aggregated} was calculated using the following formula:

$$s_{aggregated}^2 = \frac{1}{k^2} \sum s_i^2$$

with si the variance associated with the mean value i being averaged and k the total number of values being averages.

Calculation of effect size

All MPA outcomes (except in two cases; see below) were modeled as the natural logarithm response ratio of the indicator measured in the MPA and outside of the MPA.⁸⁸ For each study *i*, the MPA effect size (InRR_i) was calculated as the log ratio difference of the mean value of the indicator in $(\overline{X}_{MPA,i})$ and outside of $(\overline{X}_{control,i})$ the MPA (CI design) or after $(\overline{X}_{after,i})$ and before $(\overline{X}_{before,i})$ MPA establishment (BA design):

$$\ln RR_{i} = \ln \left(\frac{X_{MPA_{i}}}{\overline{X}_{control}} \right) \text{ for CI designs.}$$

$$\ln RR_{i} = \ln \left(\frac{\overline{X}_{MPA_{i}}}{\overline{X}_{actor}} \right) \text{ for BA designs.}$$

In the case of BACI design, we calculated the effect size as the log ratio of the ratio of the mean indicator value after MPA establishment in $(\overline{X}_{after MPA,i})$ and outside of $(\overline{X}_{after \ control,j})$ MPAs and before MPA establishment in $(\overline{X}_{before \ MPA,j})$ and outside of $(\overline{X}_{before \ control,j})$ MPAs:

$$\ln RR_{i} = \ln \left(\frac{\overline{X}_{after MPA,i} / \overline{X}_{before MPA,i}}{\overline{X}_{after control,i} / \overline{X}_{before control,i}} \right)$$

Positive In RR_i indicated that the pathway investigated had a higher mean value in the MPA than in the control site. Effect sizes were calculated so that a higher mean value was equivalent to a positive MPA outcome. As a result, the effect size used for the stability pathway was calculated by taking the natural logarithm of the opposite of the ratios detailed above. This is because the indicator used to measure stability (coefficient of variation) is inversely proportional to stability; i.e., the greater the coefficient of variation, the lower the stability.

The within-study variance vi associated with each effect sizes was calculated as follows:

$$V_{i} = \frac{s_{MPA,i}^{2}}{n_{MPA,i} * \overline{X}_{MPA,i}} + \frac{s_{control,i}^{2}}{n_{control,i} * \overline{X}_{control,i}} \text{ for CI designs,}$$

where $s_{control,i}^2$ and $s_{MPA,i}^2$ are the variances of $\overline{X}_{control,i}$ and $\overline{X}_{MPA,i}$, respectively, ---- are the associated sample sizes:

and
$$n_{control,i}$$
 and $n_{MPA,i}$ are the associated sample sizes

$$V_{j} = \frac{\sigma_{before MPA,j}}{n_{before MPA,i} + \overline{X}_{before MPA,j}^{2}} + \frac{\sigma_{after MPA,j}}{n_{after MPA,j} + \overline{X}_{after MPA,j}^{2}}$$
 for BA designs,

where $s_{before MPA,i}^2$ and $s_{after,i}^2$ are the variances of $\overline{X}_{before MPA,i}$ and $\overline{X}_{after MPA,i}$,

respectively, and
$$n_{before MPA_j}$$
 and $n_{affer MPA_j}$ are the associated sample sizes;
 $v_j = \frac{s_{before contral_j}^2}{s_{before MPA_j}^2} + \frac{s_{before MPA_j}^2}{s_{before MPA_j}^2} + \frac{s_{$

 $\frac{s_{before MPA,i}}{n_{before MPA,i} \overline{X}_{before MPA,i}^2}$ for BACI designs,

where $s_{before MPA,i}^2$, $s_{after,i}^2$, $s_{before control,i}^2$ and $s_{after control,i}^2$ are the variances of $\overline{X}_{before MPA,i}$, $\overline{X}_{after MPA,i}$, $\overline{X}_{before control,i}$, and $\overline{X}_{after control,i}$, respectively, and nbefore MPA, i, nafter MPA, i, nbefore control, i, and nafter control, i are the associated sample

sizes

The effect size used for sediment accretion (coastal protection indicator) and pH buffering were not calculated as a log ratio but as an Euclidean difference. This was done because values of accretion were sometimes positive for the MPA and negative for the control (because of erosion), which did not allow us to calculate log ratios, and because variations in pH are best measured through differences. As a result, the MPA effect size di and variance vi for sediment accretion and pH buffering was calculated as follows:

$$d_i = \overline{X}_{MPA,i} - \overline{X}_{control,i}$$
$$V_i = S^2 + S^2_{control,i}$$

Parametric data analysis

We used a weighted random-effects model to quantify the effect of MPAs on each indicator. Effect sizes were weighted accounting for the within- and among-study variance components. 89 Models were fitted using the {metafor)⁹⁰ package under R Core Team (2021).⁸⁶ Model heterogeneity, residual heterogeneity, degrees of freedom, and p values associated with each model tested are detailed in Table S4. MPAs were considered to have a significant effect of an indicator when the 95% confidence interval calculated by the model did not overlap zero.

Sensitivity analysis

To test the robustness of our meta-analysis results, we carried out a sensitivity analysis to detect (1) the presence of publication bias and of outliers using visual observation of funnel plots⁹¹ (Figures S5.1-S5.6), (2) the sensitivity of our results to publication bias using Rosenthal's fail-safe number (N_{fs}) ,⁹ whether a different outcome could be obtained when correcting for publication bias using Duval and Tweedie's trim and fill test,93,94 and (4) the impact of outlier removal on our results. For meta-analyses that did not have a significant result in the first place (Shannon index, recruitment and recovery), only the trim and fill test was performed.

Rosenthal's N_{fs}

Rosenthal's N_{fs} is an estimation of the number of additional non-significant effect sizes required for a significant meta-analysis result to become non-significant. This allows us to check the sensitivity of results to uncaptured studies. This risk is estimated to be high when N_{fs} is below the 5n+10, with n the number of data points in the meta-analysis.⁹² This was the case for only one meta-analysis result (resistance), which was the only meta-analysis showing a negative effect of MPAs on resilience attributes (Table S4).

Trim and fill test

Nine of 12 meta-analyses showed robust results of the trim and fill test (Table S4). The trim and fill test detected two potentially unstable meta-analysis results: C sequestration by sediments and sediment accretion by blue C ecosystems.

Impact of outliers on results

Outlying effect sizes were identified through visual observation of funnel plots⁹¹ (Figures S5.1–S5.6). The weight attributed to each effect size by the random-effects models was checked using forest plots (Figures S5.1-S5.6). This was done to make sure no data point was overrepresented by a much higher weight than those of the other data points. This was not the case for any of our meta-analyses. The studies corresponding to outlying points were scrutinized for factors that could explain the extreme values found. Because no flaws or marked differences in experimental design of these studies were



found, no points were excluded from our meta-analyses. One-by-one removal of outliers did not affect any of our meta-analysis results except for the resistance indicator, which became non-significant (Table S4).

Non-parametric data analysis

The indicators associated with genetic diversity and variability (allelic richness and coefficient of variation, respectively) were not associated with errors in the literature because of how these indicators are calculated. To provide quantitative results from those indicators, we performed non-parametric analyses using bootstrapping. Our bootstrapping analysis was weighted using the sample size associated with each value. MPAs were considered to have a significant effect on an indicator when the 95% confidence interval calculated through bootstrapping (n = 1,000 iterations) did not overlap zero.

Influence of moderators

Influence of moderators was investigated for pathways for which MPAs had a significant positive effect. Heterogeneity tests were run to assess how features (MPA level of protection, size [log transformed], age, presence of a fully protected area, and habitat) could mediate MPA outcomes on climate pathways (Table S4). All analyses were carried out in R using the {metafor} package.⁹⁰

MPAs of poor and moderate protection levels were grouped into a "low level" of protection because sample sizes were too small to evaluate their effect separately. As a result, the effect of level of protection was assessed for full, high, and low protection.

C sequestration from fish biomass

The C sequestration pathway was investigated using keywords related to "carbon storage" or "carbon sequestration," which returned very few results for fish. However, C storage in living fish biomass is directly proportional to fish biomass, which has been extensively measured in MPA research. Because those studies were not found through our dedicated literature search for C sequestration and because many synthetic studies and meta-analyses have already quantified the effect of MPAs on fish biomass, we based the value of this effect size on previous meta-analyses.

The effect of MPAs on fish biomass was calculated from three previous meta-analyses.^{15,16,61} Those three meta-analyses were chosen because they used completely distinct datasets, were among the most cited publications for effects of MPAs on fish biomass, and, together, encompassed all types of protection levels and all oceanic regions of the world. For the effect of level of protection on fish biomass, we used values given in Lester et al.¹⁶ for the effects of low and high protection.

Capacity of habitats to provide climate benefits

Baseline capacity of habitats to provide C sequestration and coastal protection services, as well as global habitat extent, was based on values from the most recent and global available published reviews. Values and the reference of the study from which the values were extracted are detailed in Table S4.

We also performed a systematic search and meta-analysis following the methodology presented in the "qualitative data extraction" and "quantitative data extraction" sections to estimate the amount of C stored in macroalgae and the capacity of mangroves, seagrass, and macroalgae to provide acidity buffering. This was done because we found no reviews that provided these values. The search string and date of the systematic search as well as the indicators selected were the same as those specified for the C sequestration and acidity buffering pathway (Data S1; see "data and code availability" section). Studies included in the meta-analysis on pH buffering capacity of marine habitats and on C storage capacity of macroalgae can be found in Data S3 and Data S4, respectively (see "data and code availability" section).

Resistance, recovery, and adaptation potential values

Observed resistance and recovery were calculated through meta-analysis according to the methodology presented in the "qualitative data extraction" and "quantitative data extraction" sections. The search string and date of the systematic search as well as the indicators selected are detailed in Data S1 (see "data and code availability" section).

Expected ecological adaptation was calculated as the mean value of all effect sizes contributing to ecological adaptation, i.e., allelic richness (genetic diversity), species richness and Shannon index (biodiversity), temporal stability (stability), and reproductive output and recruitment (reproduction). The confidence interval was calculated by bootstrapping, using n = 1,000 iterations.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at https://doi.org/10.1016/j. oneear.2022.09.002.

ACKNOWLEDGMENTS

Support for this study was provided by Pew Bertarelli Ocean Legacy. The views expressed are those of the authors and do not necessarily reflect the views of Pew Bertarelli Ocean Legacy.

AUTHOR CONTRIBUTIONS

This study was conceptualized by J.C.. The methodology for this was developed by J.C. and J.J. The systematic reviews and meta-analyses were carried out by J.J. with help from C.L.C. and M.L.G. under the supervision of J.C.. The original draft was written by J.J. and J.C., and reviews were performed by J.J., J.C., and R.B.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

- IPCC (2022). Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. <u>https://www.ipcc. ch/report/ar6/wg2/</u>.
- IPCC (2019). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. https://www.ipcc.ch/srocc/.
- Gallo, N.D., Victor, D.G., and Levin, L.A. (2017). Ocean commitments under the Paris agreement. Nat. Clim. Chang. 7, 833–838. https://doi.org/10. 1038/nclimate3422.
- Hoegh-Guldberg, O., and Bruno, J.F. (2010). The impact of climate change on the world's marine ecosystems. Science 328, 1523–1528. https://doi. org/10.1126/science.1189930.
- Gattuso, J.-P., Magnan, A., Billé, R., Cheung, W.W.L., Howes, E.L., Joos, F., Allemand, D., Bopp, L., Cooley, S.R., Eakin, C.M., et al. (2015). Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios. Science 349, aac4722. https://doi.org/10.1126/science.aac4722.
- Thiault, L., Mora, C., Cinner, J.E., Cheung, W.W.L., Graham, N.A.J., Januchowski-Hartley, F.A., Mouillot, D., Sumaila, U.R., and Claudet, J. (2019). Escaping the perfect storm of simultaneous climate change impacts on agriculture and marine fisheries. Sci. Adv. 5, eaaw9976. https://doi.org/ 10.1126/sciadv.aaw9976.
- Gattuso, J.-P., Magnan, A.K., Bopp, L., Cheung, W.W.L., Duarte, C.M., Hinkel, J., Mcleod, E., Micheli, F., Oschlies, A., Williamson, P., et al. (2018). Ocean solutions to address climate change and its effects on marine ecosystems. Front. Mar. Sci. 5, 337. https://doi.org/10.3389/fmars. 2018.00337.
- Reimer, J.M., Devillers, R., and Claudet, J. (2020). Benefits and gaps in areabased management tools for the ocean Sustainable Development Goal. Nat. Sustain. 4, 349–357. https://doi.org/10.1038/s41893-020-00659-2.
- Grorud-Colvert, K., Sullivan-Stack, J., Roberts, C., Constant, V., Horta e Costa, B., Pike, E.P., Kingston, N., Laffoley, D., Sala, E., Claudet, J., et al. (2021). The MPA Guide: a framework to achieve global goals for the ocean. Science 373, eabf0861. https://doi.org/10.1126/science.abf0861.
- Roberts, C.M., O'Leary, B.C., McCauley, D.J., Cury, P.M., Duarte, C.M., Lubchenco, J., Pauly, D., Sáenz-Arroyo, A., Sumaila, U.R., Wilson, R.W., et al. (2017). Marine reserves can mitigate and promote adaptation to climate change. Proc. Natl. Acad. Sci. USA *114*, 6167–6175. https://doi. org/10.1073/pnas.1701262114.
- 11. Hilborn, R. (2018). Are MPAs effective? ICES J. Mar. Sci. 75, 1160–1162. https://doi.org/10.1093/icesjms/fsx068.
- Bates, A.E., Cooke, R.S., Duncan, M.I., Edgar, G.J., Bruno, J.F., Benedetti-Cecchi, L., Côté, I.M., Lefcheck, J.S., Costello, M.J., Barrett, N., et al. (2019). Climate resilience in marine protected areas and the 'Protection Paradox. Biol. Conserv. 236, 305–314. https://doi.org/10.1016/j.biocon.2019.05.005.
- Epstein, G., Middelburg, J.J., Hawkins, J.P., Norris, C.R., and Roberts, C.M. (2022). The impact of mobile demersal fishing on carbon storage in



seabed sediments. Glob. Chang. Biol. 28, 2875–2894. https://doi.org/10. 1111/gcb.16105.

- Pendleton, L.H., Ahmadia, G.N., Browman, H.I., Thurstan, R.H., Kaplan, D.M., and Bartolino, V. (2018). Debating the effectiveness of marine protected areas. ICES J. Mar. Sci. 75, 1156–1159. https://doi.org/10.1093/ icesjms/fsx154.
- Zupan, M., Fragkopoulou, E., Claudet, J., Erzini, K., Horta e Costa, B., and Gonçalves, E.J. (2018). Marine partially protected areas: drivers of ecological effectiveness. Front. Ecol. Environ. 16, 381–387. https://doi.org/10. 1002/fee.1934.
- Lester, S., Halpern, B., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B., Gaines, S., Airamé, S., and Warner, R. (2009). Biological effects within notake marine reserves: a global synthesis. Mar. Ecol. Prog. Ser. 384, 33–46. https://doi.org/10.3354/meps08029.
- Giakoumi, S., Scianna, C., Plass-Johnson, J., Micheli, F., Grorud-Colvert, K., Thiriet, P., Claudet, J., Di Carlo, G., Di Franco, A., Gaines, S.D., et al. (2017). Ecological effects of full and partial protection in the crowded Mediterranean Sea: a regional meta-analysis. Sci. Rep. 7, 8940. https://doi. org/10.1038/s41598-017-08850-w.
- O'Leary, J.K., Micheli, F., Airoldi, L., Boch, C., De Leo, G., Elahi, R., Ferretti, F., Graham, N.A.J., Litvin, S.Y., Low, N.H., et al. (2017). The resilience of marine ecosystems to climatic disturbances. BioScience 67, 208–220. https://doi.org/10.1093/biosci/biw161.
- Miller, D.D., Ota, Y., Sumaila, U.R., Cisneros-Montemayor, A.M., and Cheung, W.W.L. (2018). Adaptation strategies to climate change in marine systems. Glob. Chang. Biol. 24, e1–e14. https://doi.org/10.1111/gcb.13829.
- Bernhardt, J.R., and Leslie, H.M. (2013). Resilience to climate change in coastal marine ecosystems. Ann. Rev. Mar. Sci 5, 371–392. https://doi. org/10.1146/annurev-marine-121211-172411.
- Freedman, R.M., Brown, J.A., Caldow, C., and Caselle, J.E. (2020). Marine protected areas do not prevent marine heatwave-induced fish community structure changes in a temperate transition zone. Sci. Rep. 10, 21081. https://doi.org/10.1038/s41598-020-77885-3.
- Mellin, C., Aaron MacNeil, M., Cheal, A.J., Emslie, M.J., and Julian Caley, M. (2016). Marine protected areas increase resilience among coral reef communities. Ecol. Lett. 19, 629–637. https://doi.org/10.1111/ele.12598.
- Bruno, J.F., Côté, I.M., and Toth, L.T. (2019). Climate change, coral loss, and the curious case of the parrotfish paradigm: why don't marine protected areas improve reef resilience? Ann. Rev. Mar. Sci 11, 307–334. https://doi.org/10.1146/annurev-marine-010318-095300.
- Di Lorenzo, M., Guidetti, P., Di Franco, A., Calò, A., and Claudet, J. (2020). Assessing spillover from marine protected areas and its drivers: a meta-analytical approach. Fish Fish. 21, 906–915. https://doi.org/10. 1111/faf.12469.
- Ban, N.C., Gurney, G.G., Marshall, N.A., Whitney, C.K., Mills, M., Gelcich, S., Bennett, N.J., Meehan, M.C., Butler, C., Ban, S., et al. (2019). Well-being outcomes of marine protected areas. Nat. Sustain. 2, 524–532. https:// doi.org/10.1038/s41893-019-0306-2.
- Hilborn, R. (2016). Policy: marine biodiversity needs more than protection. Nature 535, 224–226. https://doi.org/10.1038/535224a.
- Cinner, J., and Huchery, C. (2014). A comparison of social outcomes associated with different fisheries Co-management institutions: fisheries co-management. Conserv. Lett. 7, 224–232. https://doi.org/10.1111/ conl.12057.
- Cinner, J.E., Adger, W.N., Allison, E.H., Barnes, M.L., Brown, K., Cohen, P.J., Gelcich, S., Hicks, C.C., Hughes, T.P., Lau, J., et al. (2018). Building adaptive capacity to climate change in tropical coastal communities. Nat. Clim. Chang. 8, 117–123. https://doi.org/10.1038/s41558-017-0065-x.
- Mora, C., Wei, C.-L., Rollo, A., Amaro, T., Baco, A.R., Billett, D., Bopp, L., Chen, Q., Collier, M., Danovaro, R., et al. (2013). Biotic and human vulnerability to projected changes in ocean biogeochemistry over the 21st century. PLoS Biol. *11*, e1001682. https://doi.org/10.1371/journal.pbio.1001682.
- Cinner, J.E., Zamborain-Mason, J., Gurney, G.G., Graham, N.A.J., Mac-Neil, M.A., Hoey, A.S., Mora, C., Villéger, S., Maire, E., McClanahan, T.R., et al. (2020). Meeting fisheries, ecosystem function, and biodiversity goals in a human-dominated world. Science 368, 307–311. https://doi. org/10.1126/science.aax9412.
- Mariani, G., Cheung, W.W.L., Lyet, A., Sala, E., Mayorga, J., Velez, L., Gaines, S.D., Dejean, T., Troussellier, M., and Mouillot, D. (2020). Let more big fish sink: fisheries prevent blue carbon sequestration—half in unprofitable areas. Sci. Adv. 6, eabb4848. https://doi.org/10.1126/sciadv. abb4848.
- Mascia, M.B., Claus, C.A., and Naidoo, R. (2010). Impacts of marine protected areas on fishing communities. Conserv. Biol. 24, 1424–1429. https://doi.org/10.1111/j.1523-1739.2010.01523.x.



- Wittmann, A.C., and Pörtner, H.O. (2013). Sensitivities of extant animal taxa to ocean acidification. Nat. Clim. Chang. 3, 995–1001. https://doi. org/10.1038/nclimate1982.
- Leung, J.Y.S., Zhang, S., and Connell, S.D. (2022). Is ocean acidification really a threat to marine calcifiers? A systematic review and meta-analysis of 980+ studies spanning two decades. Small 18, 2107407. https://doi. org/10.1002/smll.202107407.
- Turnbull, J.W., Johnston, E.L., and Clark, G.F. (2021). Evaluating the social and ecological effectiveness of partially protected marine areas. Conserv. Biol. 35, 921–932. https://doi.org/10.1111/cobi.13677.
- 36. Groupe d'experts intergouvernemental sur l'évolution du climat (2014). 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands: Methodological Guidance on Lands with Wet and Drained Soils, and Constructed Wetlands for Wastewater Treatment (Intergovernmental Panel on Climate Change).
- Krause-Jensen, D., Lavery, P., Serrano, O., Marbà, N., Masque, P., and Duarte, C.M. (2018). Sequestration of macroalgal carbon: the elephant in the Blue Carbon room. Biol. Lett. *14*, 20180236. https://doi.org/10. 1098/rsbl.2018.0236.
- Griscom, B.W., Adams, J., Ellis, P.W., Houghton, R.A., Lomax, G., Miteva, D.A., Schlesinger, W.H., Shoch, D., Siikamäki, J.V., Smith, P., et al. (2017). Natural climate solutions. Proc. Natl. Acad. Sci. USA *114*, 11645–11650. https://doi.org/10.1073/pnas.1710465114.
- Yang, W., Xia, L., Zhu, Z., Jiang, L., Cheng, X., and An, S. (2019). Shift in soil organic carbon and nitrogen pools in different reclaimed lands following intensive coastal reclamation on the coasts of eastern China. Sci. Rep. 9, 5921. https://doi.org/10.1038/s41598-019-42048-6.
- Saba, G.K., Burd, A.B., Dunne, J.P., Hernández-León, S., Martin, A.H., Rose, K.A., Salisbury, J., Steinberg, D.K., Trueman, C.N., Wilson, R.W., et al. (2021). Toward a better understanding of fish-based contribution to ocean carbon flux. Limnol. Oceanogr. 66, 1639–1664. https://doi.org/ 10.1002/Ino.11709.
- Martin, S.L., Ballance, L.T., and Groves, T. (2016). An ecosystem services perspective for the oceanic eastern tropical pacific: commercial fisheries, carbon storage, recreational fishing, and biodiversity. Front. Mar. Sci. 3. https://doi.org/10.3389/fmars.2016.00050.
- Stafford, R., Boakes, Z., Hall, A.E., and Jones, G.C.A. (2022). The role of predator removal by fishing on ocean carbon dynamics. Anthr. Sci. 1, 204–210. https://doi.org/10.1007/s44177-021-00005-x.
- Ortega, A., Geraldi, N.R., Alam, I., Kamau, A.A., Acinas, S.G., Logares, R., Gasol, J.M., Massana, R., Krause-Jensen, D., and Duarte, C.M. (2019). Important contribution of macroalgae to oceanic carbon sequestration. Nat. Geosci. *12*, 748–754. https://doi.org/10.1038/s41561-019-0421-8.
- 44. Cyronak, T., Andersson, A.J., D'Angelo, S., Bresnahan, P., Davidson, C., Griffin, A., Kindeberg, T., Pennise, J., Takeshita, Y., and White, M. (2018). Short-term spatial and temporal carbonate chemistry variability in two contrasting seagrass meadows: implications for pH buffering capacities. Estuar. Coast *41*, 1282–1296. https://doi.org/10.1007/s12237-017-0356-5.
- Claudet, J., Osenberg, C.W., Benedetti-Cecchi, L., Domenici, P., García-Charton, J.A., Pérez-Ruzafa, A., Badalamenti, F., Bayle-Sempere, J., Brito, A., Bulleri, F., et al. (2008). Marine reserves: size and age do matter. Ecol. Lett. *11*, 481–489. https://doi.org/10.1111/j.1461-0248.2008. 01166.x.
- Duffy, J.E., Lefcheck, J.S., Stuart-Smith, R.D., Navarrete, S.A., and Edgar, G.J. (2016). Biodiversity enhances reef fish biomass and resistance to climate change. Proc. Natl. Acad. Sci. USA *113*, 6230–6235. https://doi. org/10.1073/pnas.1524465113.
- Yachi, S., and Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. Proc. Natl. Acad. Sci. USA 96, 1463–1468. https://doi.org/10.1073/pnas.96.4.1463.
- Hixon, M.A., Johnson, D.W., and Sogard, S.M. (2014). BOFFFFs: on the importance of conserving old-growth age structure in fishery populations. ICES J. Mar. Sci. 71, 2171–2185. https://doi.org/10.1093/icesjms/fst200.
- Planes, S., Galzin, R., Rubies, A.G., Goñi, R., Harmelin, J.G., Diréach, L.L., Lenfant, P., and Quetglas, A. (2000). Effects of marine protected areas on recruitment processes with special reference to Mediterranean littoral ecosystems. Environ. Conserv. 27, 126–143. https://doi.org/10.1017/ S0376892900000175.
- Micheli, F., Saenz-Arroyo, A., Greenley, A., Vazquez, L., Espinoza Montes, J.A., Rossetto, M., and De Leo, G.A. (2012). Evidence that marine reserves enhance resilience to climatic impacts. PLoS One 7, e40832. https://doi. org/10.1371/journal.pone.0040832.
- 51. Ferrario, F., Beck, M.W., Storlazzi, C.D., Micheli, F., Shepard, C.C., and Airoldi, L. (2014). The effectiveness of coral reefs for coastal hazard risk

reduction and adaptation. Nat. Commun. 5, 3794. https://doi.org/10.1038/ ncomms4794.

- Shepard, C.C., Crain, C.M., and Beck, M.W. (2011). The protective role of coastal marshes: a systematic review and meta-analysis. PLoS One 6, e27374. https://doi.org/10.1371/journal.pone.0027374.
- Wennerström, L., Jansson, E., and Laikre, L. (2017). Baltic Sea genetic biodiversity: current knowledge relating to conservation management. Aquat. Conserv. 27, 1069–1090. https://doi.org/10.1002/aqc.2771.
- Hadi, S., Andayani, N., Muttaqin, E., Simeon, B.M., Ichsan, M., Subhan, B., and Madduppa, H. (2020). Genetic connectivity of the scalloped hammerhead shark Sphyma lewini across Indonesia and the Western Indian Ocean. PLoS One 15, e0230763. https://doi.org/10.1371/journal.pone. 0230763.
- 55. Carr, M.H., Robinson, S.P., Wahle, C., Davis, G., Kroll, S., Murray, S., Schumacker, E.J., and Williams, M. (2017). The central importance of ecological spatial connectivity to effective coastal marine protected areas and to meeting the challenges of climate change in the marine environment. Aquat. Conserv. 27, 6–29. https://doi.org/10.1002/aqc.2800.
- Vandeperre, F., Higgins, R.M., Sánchez-Meca, J., Maynou, F., Goñi, R., Martín-Sosa, P., Pérez-Ruzafa, A., Afonso, P., Bertocci, I., Crec'hriou, R., et al. (2011). Effects of no-take area size and age of marine protected areas on fisheries yields: a meta-analytical approach. Fish Fish. 12, 412–426. https://doi.org/10.1111/j.1467-2979.2010.00401.x.
- van de Geer, C., Mills, M., Adams, V.M., Pressey, R.L., and McPhee, D. (2013). Impacts of the moreton bay marine park rezoning on commercial fishermen. Mar. Pol. 39, 248–256. https://doi.org/10.1016/j.marpol.2012. 11.006.
- Yang, Y.-C., Wang, H.-Z., and Chang, S.-K. (2013). Social dimensions in the success of a marine protected area: a case in a Taiwan fishing community. Coast. Manag. 41, 161–171. https://doi.org/10.1080/08920753.2013. 768519.
- Gollan, N., and Barclay, K. (2020). It's not just about fish": assessing the social impacts of marine protected areas on the wellbeing of coastal communities in New South Wales. PLoS One 15, e0244605. https://doi.org/10. 1371/journal.pone.0244605.
- Villegas-Ríos, D., Claudet, J., Freitas, C., Moland, E., Thorbjørnsen, S.H., Alonso-Fernández, A., and Olsen, E.M. (2021). Time at risk: individual spatial behaviour drives effectiveness of marine protected areas and fitness. Biol. Conserv. 263, 109333. https://doi.org/10.1016/j.biocon. 2021.109333.
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S., Becerro, M.A., Bernard, A.T.F., Berkhout, J., et al. (2014). Global conservation outcomes depend on marine protected areas with five key features. Nature 506, 216–220. https://doi.org/10.1038/ nature13022.
- Roff, G., and Mumby, P.J. (2012). Global disparity in the resilience of coral reefs. Trends Ecol. Evol. 27, 404–413. https://doi.org/10.1016/j.tree.2012. 04.007.
- Hoegh-Guldberg, O., Poloczanska, E.S., Skirving, W., and Dove, S. (2017). Coral reef ecosystems under climate change and ocean acidification. Front. Mar. Sci. 4. https://doi.org/10.3389/fmars.2017.00158.
- 64. IPBES (2019). Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. S. Díaz, J. Settele, E.S. Brondízio, H.T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K.A. Brauman, S.H.M. Butchart, et al., eds. (IPBES Secretariat). https:// doi.org/10.5281/zenodo.3553579.
- Cheung, W.W.L., Jones, M.C., Lam, V.W.Y., D Miller, D., Ota, Y., Teh, L., and Sumaila, U.R. (2017). Transform high seas management to build climate resilience in marine seafood supply. Fish Fish. 18, 254–263. https://doi.org/10.1111/faf.12177.
- Hopf, J.K., Jones, G.P., Williamson, D.H., and Connolly, S.R. (2019). Marine reserves stabilize fish populations and fisheries yields in disturbed coral reef systems. Ecol. Appl. 29, e01905. https://doi.org/10.1002/ eap.1905.
- 67. Emslie, M.J., Logan, M., Williamson, D.H., Ayling, A.M., MacNeil, M.A., Ceccarelli, D., Cheal, A.J., Evans, R.D., Johns, K.A., Jonker, M.J., et al. (2015). Expectations and outcomes of reserve network performance following Re-zoning of the great barrier reef marine park. Curr. Biol. 25, 983–992. https://doi.org/10.1016/j.cub.2015.01.073.
- Tittensor, D.P., Beger, M., Boerder, K., Boyce, D.G., Cavanagh, R.D., Cosandey-Godin, A., Crespo, G.O., Dunn, D.C., Ghiffary, W., Grant, S.M., et al. (2019). Integrating climate adaptation and biodiversity conservation in the global ocean. Sci. Adv. 5, eaay9969. https://doi.org/10.1126/sciadv. aay9969.



- Wilson, K.L., Tittensor, D.P., Worm, B., and Lotze, H.K. (2020). Incorporating climate change adaptation into marine protected area planning. Glob. Chang. Biol. 26, 3251–3267. https://doi.org/10.1111/gcb.15094.
- Claudet, J., Loiseau, C., and Pebayle, A. (2021). Critical gaps in the protection of the second largest exclusive economic zone in the world. Mar. Pol. 124, 104379. https://doi.org/10.1016/j.marpol.2020.104379.
- Claudet, J., Loiseau, C., Sostres, M., and Zupan, M. (2020). Underprotected marine protected areas in a global biodiversity hotspot. One Earth 2, 380–384. https://doi.org/10.1016/j.oneear.2020.03.008.
- Roessger, J., Claudet, J., and Horta e Costa, B. (2022). Turning the tide on protection illusions: The underprotected MPAs of the 'OSPAR Regional Sea Convention. Mar. Policy 142, 105109. https://doi.org/10.1016/j. marpol.2022.105109.
- Weeks, R., and Jupiter, S.D. (2013). Adaptive comanagement of a marine protected area network in Fiji. Conserv. Biol. 27, 1234–1244. https://doi. org/10.1111/cobi.12153.
- Christie, P., Bennett, N.J., Gray, N.J., 'Aulani Wilhelm, T., Lewis, N., Parks, J., Ban, N.C., Gruby, R.L., Gordon, L., Day, J., et al. (2017). Why people matter in ocean governance: incorporating human dimensions into large-scale marine protected areas. Mar. Pol. 84, 273–284. https://doi. org/10.1016/j.marpol.2017.08.002.
- Chaigneau, T., and Brown, K. (2016). Challenging the win-win discourse on conservation and development: analyzing support for marine protected areas. Ecol. Soc. 21, art36. https://doi.org/10.5751/ES-08204-210136.
- Di Franco, A., Thiriet, P., Di Carlo, G., Dimitriadis, C., Francour, P., Gutiérrez, N.L., Jeudy de Grissac, A., Koutsoubas, D., Milazzo, M., Otero, M.D.M., et al. (2016). Five key attributes can increase marine protected areas performance for small-scale fisheries management. Sci. Rep. 6, 38135. https://doi.org/10.1038/srep38135.
- 77. Di Franco, A., Hogg, K.E., Calò, A., Bennett, N.J., Sévin-Allouet, M.A., Esparza Alaminos, O., Lang, M., Koutsoubas, D., Prvan, M., Santarossa, L., et al. (2020). Improving marine protected area governance through collaboration and co-production. J. Environ. Manage. 269, 110757. https://doi.org/10.1016/j.jenvman.2020.110757.
- Bennett, N.J., and Dearden, P. (2014). Why local people do not support conservation: community perceptions of marine protected area livelihood impacts, governance and management in Thailand. Mar. Pol. 44, 107–116. https://doi.org/10.1016/j.marpol.2013.08.017.
- Voyer, M., Gladstone, W., and Goodall, H. (2015). Obtaining a social licence for MPAs – influences on social acceptability. Mar. Pol. 51, 260–266. https://doi.org/10.1016/j.marpol.2014.09.004.
- Voyer, M., Gladstone, W., and Goodall, H. (2014). Understanding marine park opposition: the relationship between social impacts, environmental knowledge and motivation to fish. Aquat. Conserv. 24, 441–462. https:// doi.org/10.1002/aqc.2363.
- Macreadie, P.I., Costa, M.D.P., Atwood, T.B., Friess, D.A., Kelleway, J.J., Kennedy, H., Lovelock, C.E., Serrano, O., and Duarte, C.M. (2021). Blue carbon as a natural climate solution. Nat. Rev. Earth Environ. 2, 826–839. https://doi.org/10.1038/s43017-021-00224-1.
- Cisneros-Montemayor, A.M., Ducros, A.K., Bennett, N.J., Fusco, L.M., Hessing-Lewis, M., Singh, G.G., and Klain, S.C. (2022). Agreements and benefits in emerging ocean sectors: are we moving towards an equitable Blue Economy? Ocean Coast Manag. 220, 106097. https://doi.org/10. 1016/j.ocecoaman.2022.106097.
- Lecerf, M., Herr, D., Thomas, T., Elverum, C., Delrieu, E., and Picourt, L. (2021). Coastal and marine ecosystems as nature-based solutions in new or updated nationally determined contributions (ocean & climate platform, conservation international, IUCN, GIZ, rare, the nature conservancy and WWF). https://ocean-climate.org/wp-content/uploads/2021/ 06/coastal-and-marine-ecosystem-2806.pdf.
- O'Connor, J.J., Fest, B.J., Sievers, M., and Swearer, S.E. (2020). Impacts of land management practices on blue carbon stocks and greenhouse gas fluxes in coastal ecosystems—a meta-analysis. Glob. Chang. Biol. 26, 1354–1366. https://doi.org/10.1111/gcb.14946.
- Sasmito, S.D., Taillardat, P., Clendenning, J.N., Cameron, C., Friess, D.A., Murdiyarso, D., and Hutley, L.B. (2019). Effect of land-use and land-cover change on mangrove blue carbon: a systematic review. Glob. Chang. Biol. 25, 4291–4302. https://doi.org/10.1111/gcb.14774.
- R Core Team (2021). R: A Language and Environment for Statistical Computing. (2021) (R Foundation for Statistical Computing).
- Horta e Costa, B., Claudet, J., Franco, G., Erzini, K., Caro, A., and Gonçalves, E.J. (2016). A regulation-based classification system for Marine Protected Areas (MPAs). Mar. Pol. 72, 192–198. https://doi.org/10.1016/ j.marpol.2016.06.021.





- Hedges, L.V., Gurevitch, J., and Curtis, P.S. (1999). The meta-analysis of response ratios in experimental ecology. Ecology 80, 1150–1156. https:// doi.org/10.2307/177062.
- Hedges, L.V., and Vevea, J.L. (1998). Fixed- and random-effects models in meta-analysis. Psychological Methods 3, 486–504. https://doi.org/10. 1037/1082-989X.3.4.486.
- Viechtbauer, W. (2019). Meta-Analysis Package for R: Package "Metafor" Version 2.4-0 (CRAN - R Project).
- Page, M.J., Sterne, J.A.C., Higgins, J.P.T., and Egger, M. (2021). Investigating and dealing with publication bias and other reporting biases in meta-analyses of health research: a review. Res. Synth. Methods 12, 248–259. https://doi.org/10.1002/jrsm.1468.
- Sampaio, E., Santos, C., Rosa, I.C., Ferreira, V., Pörtner, H.O., Duarte, C.M., Levin, L.A., and Rosa, R. (2021). Impacts of hypoxic events surpass those of future ocean warming and acidification. Nat. Ecol. Evol. 5, 311–321. https://doi.org/10.1038/s41559-020-01370-3.
- Duval, S., and Tweedie, R. (2000). Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 56, 455–463. https://doi.org/10.1111/j.0006-341X.2000.00455.x.
- Shi, L., and Lin, L. (2019). The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. Medicine 98, e15987. https://doi.org/10.1097/MD. 000000000015987.